

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 421 568 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90305367.6

(51) Int. Cl.⁵: **C07H 17/08, C07D 493/22,
A01N 43/90, A61K 31/335**

(22) Date of filing: 17.05.90

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

(30) Priority: 17.05.89 GB 8911281
22.12.89 GB 8929041(43) Date of publication of application:
10.04.91 Bulletin 91/15(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE(71) Applicant: Beecham Group p.l.c.
SB House Great West Road
Brentford Middlesex TW8 9BD(GB)(72) Inventor: Baker, Geoffrey Harold, Beecham
Pharmaceuticals
Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT18 5XQ(GB)
Inventor: Dorgan, Roderick John, Beecham**Pharmaceuticals**

Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT18 5XQ(GB)
Inventor: Morgan, David Owen, Beecham
Pharmaceuticals


Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT18 5XQ(GB)
Inventor: Shelley, Peter Robin, Beecham
Pharmaceuticals
Brockham Park

Betchworth, Surrey RH3 7AJ(GB)
Inventor: Blanchflower, Simon Edward,
Beecham Pharmaceutical
Walton Oaks, Dorking Road
Tadworth, Surrey KT20 7NT(GB)

(74) Representative: West, Vivien et al
SmithKline Beecham, Corporate Patents,
Great Burgh, Yew Tree Bottom Road
Epsom, Surrey KT18 5XQ(GB)

(54) Macrolide compounds.

(57) Novel compounds of formula (I):



(I)

wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or


optionally O-substituted oximino; R^3 is hydrogen, optionally protected hydroxy, or a group $4'-(\alpha\text{-L-oleandrosyl})-\alpha\text{-L-oleandrosyloxy}$ or $\alpha\text{-L-oleandrosyloxy}$ wherein the terminal hydroxy group is optionally protected; R^4 , R^5 , R^6 and R^7 are the same or different and each is hydrogen or an organic radical; and R^8 is an optionally substituted amino or imino group such as optionally O-substituted oxyimino, optionally N-substituted hydrazone, or optionally N-substituted semicarbazone; are useful in the treatment of helminthiasis in humans and animals.

NOVEL COMPOUNDS

The present invention relates to novel anthelmintic compounds, to processes for their production, to pharmaceutical formulations containing them, and to their use in human or veterinary medicine.


The milbemycins and avermectins are a group of macrolide antibiotics which have been prepared by the cultivation of microorganisms and are described in inter alia GB-A-1,390,336, J. Antibiotics 29 (3), 76-14 to 76-16 and 29 (6), 76-35 to 76-42, GB-A-2 170 499, EP-A-O 073 660 and EP-A-0 204 421. They have anthelmintic activity. Further anthelmintically active milbemycins and avermectins are described in GB-A-2 176 180, EP-A-0 212 867, EP-A-0 237 339, EP-A-0 241 146, EP-A-0 214 731, EP-A-0 194 125, EP-A-0 170,006, and US-A-4,285,963.

The compounds disclosed in the above references include compounds of formula (A):



(A)


wherein R^a is methoxy or hydroxy, R^b is hydrogen, R^c is hydrogen, pentanoyloxy, heptanoyloxy, or 2-methylhexanoyloxy, and R^d is methyl or ethyl, with the proviso that when R^c is hydrogen, R^a is methoxy; or R^a is methoxy or hydroxy, R^b is hydrogen, R^c is 2-methylbutanoyloxy, 2,4-dimethylpent-2-enoyloxy, or 2,4-dimethylpentanoyloxy, and R^d is methyl or ethyl, with the proviso that when R^d is ethyl, R^a is hydroxy and R^c is 2,4-dimethylpentanoyloxy; or R^a is methoxy or hydroxy, R^b is the group of formula:



(4'-(α -L-oleandrosyl)- α -L-oleandrosyloxy), R^c is hydroxy, and R^d is 1-methyl propyl.
EP-A-O 254 583 (USN 076,274) describes compounds of formula (B):

5

10



(B)

15


wherein R^e is hydrogen or E 2-methyl 2-butenyloxy, and R^f is methoxy or hydroxy, with the proviso that when R^e is E 2-methyl 2-butenyloxy, R^f is methoxy.

EP-A-0325462 (USSN 299,933) describes compounds of formula (C):

20

25

30




(C)

35

wherein R^g and R^h are as set out in the following table:

40

45

Compound	R^g	R^h
VM48130	$O-CO-CH(CH_3)_2$	H
VM48633	H	$O-CO-CH=C(CH_3)_2$
VM47704	H	$O-CO-CH_2-CH(CH_3)_2$
VM48642	H	$O-CO-CH_2$ 


50

EP-A-0 288 205 (USSN 183,581) discloses compounds of formula (D):

55

5

10



(D)


15 wherein R^i is hydroxy or methoxy and R^j and R^k are the same or different and each is selected from optionally protected hydroxy, alkoxy, acyloxy, sulphonyloxy, oxo and optionally O-substituted oximino.

EP-A-0 259 779, EP-A-0 293 549, EP-A-0 307 225 and GB-A-2192630 describe compounds of formula (E):

20

25

30



(E)


35 wherein R^1 is optionally protected hydroxy or methoxy, R^m is optionally protected hydroxy, oxo, or an imino group such as optionally O-substituted oxyimino, optionally N-substituted hydrazone, or optionally N-substituted semicarbazone, R^n is methyl, ethyl or isopropyl, and the dashed line is a double bond or an epoxide group.

EP-A-0 260 536 and EP-A-0 260 537 describe compounds of formula (F):

40

45

50




(F)

55


wherein R^o is optionally protected hydroxy or methoxy, R^p is hydrogen or a sugar residue, R^q is optionally protected hydroxy, oxo, or an imino group such as optionally O-substituted oxyimino or optionally N-substituted hydrazone, and R^r is isopropyl or sec-butyl.

The absolute configuration of the compounds of formulae (A) to (F) is believed to be as follows:



We have now discovered that it is possible to prepare novel compounds from starting materials having a hydroxy substituent at the C-22 position, such as compounds of formulae (A) to (D) above and the compounds disclosed in EP-A-O 334 484, and that these compounds are useful as anthelmintically active compounds.

According to the present invention there is provided a compound of formula (I):



(I)

wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or optionally O-substituted oximino; R³ is hydrogen, optionally protected hydroxy, or a group 4'-(α -L -oleandrosyl)- α -L -oleandrosyloxy or α -L -oleandrosyloxy wherein the terminal hydroxy group is optionally protected; R⁴, R⁵, R⁶ and R⁷ are the same or different and each is hydrogen or an organic radical; and R⁸ is an optionally substituted amino or imino group such as optionally O-substituted oxyimino, optionally N-substituted hydrazone, or optionally N-substituted semicarbazone; with the proviso that the compound of formula (I) is not a compound of formula (E) or (F) above or a compound disclosed in EP-A-0 307 220. Typically R³ is in the α -configuration shown above for formulae (A) and (F). When R³ is in the β -configuration then it is preferably optionally protected hydroxy, and/or R¹ is preferably hydrogen, and/or R² is preferably methoxy or optionally protected hydroxy.










Preferably, (a) when R¹ is optionally protected hydroxy, R³ is hydrogen, and/or (b) when R⁴ is methyl and R⁵ and R⁶ are hydrogen, R⁷ does not have any of the values set out in Table I hereinbelow, and/or (c) when R⁴ is methyl and R⁵ and R⁶ are hydrogen, R⁷ does not have any of the values set out in Table II hereinbelow, and/or (d) when R² is not methoxy or optionally protected hydroxy, R¹ and R³ are both hydrogen.

Certain compounds of formula (I) wherein R⁸ is optionally protected hydroxy or oxo are also novel, and such compounds form a further aspect of the invention.

Thus, a particular aspect of the invention provides a compound of formula (I) as defined hereinabove wherein R⁸ is optionally protected by hydroxy or oxo, with the proviso that (a) when R³, R⁵ and R⁶ are hydrogen and R⁴ is methyl, then R⁷ does not have any of the values set out in Table III hereinbelow, (b) when R³ is not hydrogen, R⁴ is methyl and R⁵ and R⁶ are hydrogen, then R⁷ does not have any of the values set out in Table IV hereinbelow, (c) when R³, R⁵ and R⁶ are hydrogen and R⁴ is ethyl, then R⁷ is not (Z)-4-methyl-pent-2-en-2-yl, and (d) when R³ is not hydrogen, R⁴ is methyl, and R⁵ is hydrogen, the 25-position is not substituted in the same way as that of compound III of US-A-4,285,963, and (e) it is not a compound disclosed in EP-A-0 317 148, EP-A-0 307 220, EP-A-0 308 145, EP-A-0 350 187 or EP-A-0 335 541.

TABLE I


-CH(CH₃)₂, -CH(CH₃)C₂H₅

TABLE II

R⁹ is methyl, ethyl or isopropyl

R¹⁰ is C₁₋₇ alkyl, phenyl

or C₇₋₁₄ phenalkyl



X is halogen

R¹¹ is methyl, ethyl or isopropyl.


1-methylthioethyl, cyclobutyl, cyclopentyl, cyclohexyl, 2-methylcyclopropyl, cyclohex-3-enyl, or thien-3-yl.

TABLE III

5


 R^9 is methyl, ethyl or isopropyl

10





 R^{10} is C_{1-7} alkyl, phenyl
or C_{7-14} phenalkyl

15



X is halogen

 R^{11} is methyl, ethyl or isopropyl.

20



25





30

TABLE IV

35

 $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_3)\text{C}_2\text{H}_5$

40




45

 1-methylthioethyl, cyclobutyl, cyclopentyl,
cyclohexyl, 2-methylcyclopropyl, cyclohex-3-enyl,
or thien-3-yl.

50

preferably, when R^4 is methyl and R^5 and R^6 are hydrogen, R^7 is not a group:

55



5

wherein R⁹ is methyl, ethyl or isopropyl and R¹⁰ is alkyl, phenalkyl or phenyl.

It is also preferred that, when R⁴ is methyl and R⁵ and R⁶ are hydrogen, R⁷ is not selected from the group consisting of pent-2-yl, 3-methylbut-2-yl, hex-2-yl, pent-4-en-2-yl, 1-cyclopropylethyl, cycloheptyl, 4,4-difluorocyclohexyl, 4-methylenecyclohexyl, 3-methylcyclohexyl, cyclopenten-1-yl, cyclohexen-1-yl, tetrahydropyran-4-yl, thien-2-yl, 3-furyl, and 2-chlorothien-4-yl.

In a particular aspect, when R⁴ is methyl and R⁵ and R⁶ are hydrogen, R⁷ is not an alpha-branched C₃₋₈ alkylthioalkyl group; a C₃₋₈ cycloalkyl group optionally substituted by one C₁₋₄ alkyl group; a C₅₋₈ cycloalkenyl group; or a 3 to 6 membered sulphur containing heterocyclic ring. More especially, R⁷ is not an alpha-branched C₃₋₈ alkyl, alkenyl or alkylthioalkyl group; a C₅₋₈ cycloalkylalkyl group wherein the alkyl group is an alpha-branched C₂₋₅ alkyl group; a C₃₋₈ cycloalkyl group optionally substituted by methylene or one or more C₁₋₄ alkyl groups or halo atoms; a C₅₋₈ cycloalkenyl group; or a 3 to 6 membered oxygen or sulphur containing heterocyclic ring which may optionally be substituted by one or more halo atoms.

In a still further particular aspect, when R⁴ is methyl and R⁵ and R⁶ are hydrogen, R⁷ is not an alpha-branched C₃₋₈ alkyl, alkenyl, alkynyl, alkoxyalkyl or alkylthioalkyl group; a C₅₋₈ cycloalkylalkyl group wherein the alkyl group is an alpha-branched C₂₋₅ alkyl group; a C₃₋₈ cycloalkyl or C₅₋₈ cycloalkenyl group, either of which may optionally be substituted by methylene or one or more C₁₋₄ alkyl groups or halo atoms; or a 3 to 6 membered oxygen or sulphur containing heterocyclic ring which may be saturated, or fully or partially unsaturated and which may optionally be substituted by one or more C₁₋₄ alkyl groups or halo atoms. More especially, R⁷ is not an alpha-branched alkyl, alkenyl, alkynyl, alkoxyalkyl or alkylthioalkyl group; a cycloalkylalkyl group wherein the alkyl group is alpha-branched; a cycloalkyl or cycloalkenyl group, either of which may optionally be substituted by methylene or one or more alkyl groups or halo atoms; or an oxygen or sulphur containing heterocyclic ring which may be saturated, or fully or partially unsaturated and which may optionally be substituted by one or more alkyl groups or halo atoms.

According to a further preferred aspect, when R⁴ is ethyl and R⁵ and R⁶ are hydrogen, R⁷ is not (Z)-4-methyl-pent-2-en-2-yl.

According to a still further preferred aspect, when R⁴ is methyl and R⁵ is hydrogen, the 25-position is not substituted in the same way as that of compound III of US-A-4,285,963.

Suitable protecting groups for hydroxy include TBDMS (t-butyldimethylsilyl), and acyl. Further suitable protecting groups are described in, for example, "Protective Groups in Organic Synthesis" Theodora W. Greene, Wiley-Interscience 1981 Ch 2, 10-86.

When any of R⁴ to R⁷ is an organic radical it may advantageously be selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heterocyclyl, mono-, bi- and tri-cycloalkyl, mono-, bi- and tri-cycloalkenyl and aralkyl.

As used herein alkyl includes straight and branched C₁₋₂₀, more especially C₁₋₁₂, particularly C₁₋₆ alkyl, and alkenyl and alkynyl include straight and branched C₂₋₂₀, more especially C₂₋₁₂, particularly C₂₋₆ alkenyl and alkynyl.

When any of R⁴ to R⁷ comprises an alkyl, alkenyl or alkynyl moiety that moiety may optionally be substituted by one or more substituents selected from the group consisting of hydroxy, alkoxy, alkylthio, oxo, halogen, trifluoromethyl, and optionally substituted amino.

When used herein the term 'aryl' includes phenyl and naphthyl optionally substituted with up to five, preferably up to three, groups selected from halogen, C₁₋₆ alkyl, aryl, C₁₋₆ alkoxy, halo substituted (C₁₋₆) alkyl, hydroxy, amino, nitro, carboxy, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkoxy carbonyl-(C₁₋₆)-alkyl, C₁₋₆ alkyl carbonyloxy, or C₁₋₆ alkyl carbonyl groups.

The term 'heterocyclyl' includes saturated, unsaturated and aromatic single or fused rings comprising up to four hetero atoms in the ring selected from oxygen, nitrogen and sulphur and optionally substituted with up to three halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, halo-(C₁₋₆)-alkyl, hydroxy, amino, carboxy, C₁₋₆ alkoxy carbonyl, C₁₋₆ alkoxy carbonyl-(C₁₋₆)-alkyl, aryl or oxo groups.

Suitably the heterocyclic ring comprises from 4 to 7 ring atoms, preferably 5 to 6 atoms.

The term 'halogen' refers to fluorine, chlorine, bromine and iodine.

Particularly suitable substituents for an amino or imino group such as an oxime, hydrazone or semicarbazone group include one or more organic radicals as defined hereinabove for R⁴ to R⁷, for

example the substituents set out in EP-A-0 288 205, EP-A-0 259 779, EP-A-0 260 537, EP-A-0 260 536, GB-A-2 192 630, and EP-A-0 307 225.

Those skilled in the art will appreciate that an N-substituted imino group such as an oxime may exist as either an E or Z isomer, or as a mixture of E and Z isomers, and that an E or Z isomer may be converted to the other isomer or to a mixture of isomers by standard techniques such as acid treatment.

As used herein mono-, bi- and tri-cycloalkyl include C₃₋₂₀, especially C₃₋₁₂, more especially C₄₋₈, groups, and mono-, bi- and tri-cycloalkenyl include C₄₋₂₀, especially C₄₋₁₂, more especially C₅₋₈ groups. When any of R₄ to R₇ comprises a mono-, bi- or tri-cycloalkyl or mono-, bi- or tri-cycloalkenyl moiety, that moiety may be substituted as set out above for alkyl, alkenyl, and alkynyl, and/or by one or more substituents selected from the group consisting of methylene and alkyl. Bicyclic and tricyclic groups may be fused or bridged and are preferably attached via a carbon atom which is common to two rings.

Any two of R⁴ to R⁷ may be taken together with the carbon atom(s) to which they are attached to designate a cycloalkyl, cycloalkenyl, aryl or heterocyclyl group which may optionally be substituted as set out above.

In a preferred aspect of the invention, R¹ is hydrogen, R² is methoxy or hydroxy, and R³ is hydrogen.

In a particular aspect of the invention, R⁵ and R⁶ are hydrogen, one of R⁴ and R⁷ is hydrogen or alkyl, such as methyl, and the other of R⁴ and R⁷ is an organic radical, more especially an aryl, cycloalkyl, alkenyl, cycloalkenyl or alkyl group.

Preferably, R² is a hydroxy group.

The compound or mixture of compounds according to the invention is suitably provided in substantially pure form, for example at least 50% pure, suitably at least 60% pure, advantageously at least 75% pure, preferably at least 85% pure, more preferably at least 95% pure, especially at least 98% pure, all percentages being calculated as weight/weight. An impure or less pure form of a compound according to the invention may, for example, be used in the preparation of a more pure form of the same compound or of a related compound (for example a corresponding derivative) suitable for pharmaceutical use.

The compounds of the invention have parasitocidal properties, for example against nematodes such as *Trichostrongylus colubriformis*, and are useful for the treatment of helminthiasis in animals such as mammals, including humans and domesticated animals (including farm animals).

Accordingly the present invention also provides a compound according to the invention, for use in the treatment of the human or animal body, especially for treating endo- and ectoparasitic infestations and particularly for treating helminthiasis of domestic and farm animals.

The term helminthiasis encompasses those diseases of man and animals caused by infestation with parasitic worms such as Strongyles, Ascarids, hookworms lungworms, filarial worms and whipworms. The compound may also be used against nematodes occurring in the soil or parasitic to plants.

The compounds of the invention are also active against Arthropods. The phylum Arthropoda comprises insects -such as biting flies, lice, bugs, beetles and fleas -and arachnids - such as mites and ticks.

Thus, a broad aspect of the invention provides a method of eradicating arthropod or nematode infestations, which method comprises applying a compound according to the invention or a derivative thereof to the arthropods or nematodes or to their environment.

The present invention thus provides a pesticidal composition comprising a compound according to the invention or a derivative thereof together with a suitable carrier or excipient, such as an aerosol formulation.

The present invention also provides a pharmaceutical or veterinary composition comprising a compound according to the invention or a pharmaceutically acceptable derivative thereof together with a pharmaceutically or veterinarily acceptable carrier or excipient.

The present invention also provides a method of treatment or prophylaxis of endo- and ectoparasitic infestations, especially helminthiasis, of animals (including birds and fish), especially humans and domesticated mammals, which comprises administering an effective non-toxic amount of a compound according to the invention or a pharmaceutically acceptable derivative thereof, or a composition according to the invention, to a patient in need thereof.

The composition according to the invention may be formulated for administration in any convenient way for use in human or veterinary medicine, by analogy with other anthelmintics.

In suitable formulations the drug may be administered to animals orally (as a paste, drench, bolus, capsule or tablet), parenterally, percutaneously, as a food additive (eg granules, pellets or powder), or may be prepared as an aerosol spray formulation.

The compounds of the invention may be formulated as a mixture with each other and/or with other anthelmintics, insecticides, acaricides or other pharmacologically active substances.

Suitably the composition consists of sufficient material to provide a dose of from 0.001 to 100mg of active ingredient per kg of animal body weight per dose, more suitably 0.01 to 10mg/kg per dose.

A composition according to the invention may suitably contain from 0.1% by weight, preferably from 1.0 to 60% by weight, of the compound according to the invention (based on the total weight of the composition), depending on the method of administration.

It will be appreciated that, in some cases, it will be advisable to repeat the dosing of the infected or potentially infected human or animal with the compound of the invention according to conventional dosage regimes used with anthelmintics.


Compounds of formula (I) wherein R^8 is an imino group may be prepared by reacting the corresponding 23-Keto derivative with an amino containing compound such as an amine, hydroxylamine, hydrazine or semicarbazide to yield derivatives of the type where R^8 is optionally substituted imino, oxyimino, hydrazone or semicarbazone, respectively. Suitable processes are described in EP-A-0 259 779, EP-A-0 260 537, EP-A-0 260 536 and GB-A-2192630.

Compounds of formula (I) wherein R^8 is a substituted oxyimino group may be prepared by reacting the corresponding 23-hydroxyimino derivative with a suitable etherifying agent, for example an alkyl halide in the presence of a base such as triethylamine.

Furthermore, the 23-Keto derivative may be reduced to the alcohol using a suitable hydride reducing agent such as sodium borohydride, or L-selectride (Trade Mark Aldrich Chemical Co.).

A 23-imino derivative may be reduced to the corresponding amine using a suitable reducing agent such as sodium cyanoborohydride.


The 23-Keto derivative can be obtained by hydrating and cyclizing a compound of formula (II):



wherein R^1 to R^7 have the values set out above, and R^{18} is hydrogen or lower alkyl, more particularly C_{1-3} alkyl.

The acetylene (II) may be hydrated and cyclized to give the 23-Keto derivative by the action of an aqueous acid in the presence of a suitable mercury salt such as mercuric oxide. Other methods for the hydration of acetylenes are contained in standard texts e.g. The Chemistry of the C-C Triple Bond Ed S Patai Publ Wiley: New York and references cited therein.


The 23-keto derivative can also be obtained by cyclizing a compound of formula (V)



(V)

wherein R^1 to R^7 and R^{18} have the values set out above, and R^{20} is an optionally protected ketone such as an acetal, for example by treatment with aqueous acid followed, if required by removal of any protecting group.

Alternatively, compounds of formula (I) wherein R^8 is oxo or substituted oxyimino may be prepared by cyclizing a compound of formula (IV):




(IV)

wherein R^1 to R^8 and R^{18} have the values set out above, for example by treatment with aqueous acid.


It should be appreciated that the configuration at C21 of the compound of formula (I) will be subject to thermodynamic control (as discussed eg by P. Deslongchamps et al, Canadian J. Chem. [1981] 59, 1105); it is believed that the epimer in which the configuration at C21 is as in the naturally occurring compounds of formulae (A), (B) and (C) is normally thermodynamically favoured.

Compounds of formula (II), (IV) and (V) may be obtained from compounds of formula (III):



(III)

wherein R^1 to R^3 are as defined above, via the routes shown in Scheme I below:



Notes

(i) $MC\equiv C.CR^4R^5.CR^6R^7OP$ where M is a metalating agent such as Li or BrMg and P is a protecting group eg an acid labile protecting group such as THP (tetrahydropyranyl)


(ii) $R^{18}OH/H^+$

(iii) $MCH_2CR^8.CR^4R^5.CR^6R^7OP$


(iv) $MCH_2CR^{20}.CR^4R^5.CR^6R^7OP$

Compounds of formula (III) can be prepared as described in EP-A-O 319 142 (USSN 265,509) by cleaving the 21-22 carbon-carbon bond of a precursor having a hydroxy substituent on C22, such as compounds of formulae (A), (B), (C) and (D).


Those skilled in the art will appreciate that the process of the invention can be applied to essentially any milbemycin or avermectin having an optionally protected hydroxy (or oxo or oxime) group present at the C22 position. Furthermore, the compounds of formulae (I) and (III) can be further modified using techniques which are in themselves well known in the art and are described in, for example, Natural Products Reports 3 (2) [1986]87 et seq. and Macrolide Antibiotics [1984] Ch. 14, 553 et seq. Thus, a broad aspect of the invention provides any milbemycin or avermectin of partial formula (i) hereinbelow, as well as a process for its preparation which comprises (hydrating and) cyclizing a milbemycin or avermectin of partial formula (ii), (iii) or (iv) hereinbelow:




(i)



(ii)




(iii)



(iv)


wherein R⁴ to R⁸ and R¹⁸ have the values set out above; and/or, optionally, converting a compound of formula (i) to a different compound of formula (i) as set out hereinabove.

A further broad aspect of the invention provides a novel process for the preparation of milbemycins and avermectins of partial formula (v):




(v)


wherein R¹⁴ to R¹⁷ are the same or different and each is selected from hydrogen and an organic radical; and R¹⁹ is optionally protected hydroxy, oxo, or an optionally substituted amino or imino group such as optionally O-substituted oxyimino, or optionally N-substituted hydrazone or semicarbazone, which process comprises (hydrating and) cyclizing a corresponding milbemycin or avermectin of partial formula (vi), (vii) or (viii):



(vi)



(vii)



(viii)


wherein R^{14} to R^{19} have the values set out above; and/or, optionally, converting a compound of formula (v) to a different compound of formula (v) as set out hereinabove.

Acetylenic precursors of structure $HC \equiv C.CR^4R^5CR^6R^7OP$ may be prepared using the procedures described in EP-A-O 353 959 (USSN 387,351).

Precursors of structure $MCH_2CR^8.CR^4R^5CR^6R^7OP$ may be prepared by treating a compound of formula $CH_3CR^8.CR^4R^5CR^6R^7OP$ with a suitable metalating agent such as n -butyllithium in an inert solvent such as THF at low temperatures eg about $-70^\circ C$.

Precursors of structure $MCH_2C(OP)_2.CR^4R^5CR^6R^7OP$ may be prepared using procedures analogous to those described in EP-A-O 353 959.

The following Examples illustrate the present invention. VS 47709 is the compound of formula (III) wherein R^1 and R^3 are both hydrogen and R^2 is methoxy. VS 48927 is the compound of formula (III) wherein R^1 and R^3 are both hydrogen and R^2 is hydroxy protected with TBDMS. "Milbemycin X" represents the following structure:



Example 1

(24R, 25S)-24-methyl-23-oxo-25-phenyl-milbemycin X.

A mixture of (3S, 4S)-3-methyl-4-phenyl-4-(*t*-butyldimethylsilyloxy)-1-butyne (2.36g, 8.6 mmol) and dry THF (40 ml) was cooled to $-78^\circ C$ under nitrogen. 1.6 M *n*-Butyllithium in hexane (4.7 ml, 7.5 mmol) was added dropwise with stirring and stirring continued at $-78^\circ C$ (2h). A solution of VS 48927 prepared as described in Examples 1 to 3 of EP-A-0 319 142 (USSN 265, 509) (1.2 g, 2.2 mmol) in dry THF (4 ml) was rapidly added and the mixture stirred at $-78^\circ C$ (20 min). A solution of acetic acid (1.5 ml) in THF (10 ml) was cooled to $-78^\circ C$ and then rapidly added to the reaction mixture. After warming to R.T. 1M ammonium chloride solution (100 ml) was added and the mixture extracted with ether (3 x 40 ml). The combined ethereal extracts were washed with saturated sodium bicarbonate solution (40 ml), dried ($MgSO_4$) and evaporated.

The residue was dissolved in methanol (20 ml). Toluene-4-sulphonic acid (ca 200 mg) was added and the mixture stirred at R.T. (17h), treated with water, (100 ml) and extracted with ether (3 x 40 ml). The

combined ethereal extracts were washed with saturated sodium bicarbonate solution (40 ml), dried (MgSO_4) and evaporated. The product was purified by silica gel column chromatography with light petroleum (b.p. 40-60°C)-ethyl acetate, 3:2 as eluant (yield 0.96 g).

A portion of the above intermediate product (197 mg, 0.32 mmol) was dissolved in methanol (6 ml). A solution of mercuric oxide (7 mg, 0.03 mmol) in a mixture of concentrated sulphuric acid (0.1 ml) and water (3 ml) was added. The reaction mixture was stirred at R.T. (3 h), treated with water (50 ml) and then extracted with dichloromethane (3 x 20 ml).

The combined dichloromethane extracts were dried (MgSO_4) and evaporated. The residue was fractionated by silica gel column chromatography with light petroleum (b.p. 40-60°C)-ethyl acetate, 2:1 as eluant giving (24R,24S)-24-methyl-23-oxo-25-phenyl-milbemycin X (125 mg). ^{13}C n.m.r. (270 MHz; CDCl_3) included δ 205.70, 139.20, 128.53, 128.35, 127.50, 40.45, and 9.29ppm.

Example 2

(24S, 25S)-24-methyl-23-methoxyimino-25-phenyl-milbemycin X.




(24R, 25S)-24-methyl-23-oxo-25-phenyl-milbemycin X (203 mg, 0.34 mmol) was dissolved in methanol (20 ml). A solution of methoxylamine hydrochloride (204 mg, 2.43 mmol) in water (4 ml) was added and the mixture stirred at R.T. (4 h). The bulk of the methanol was evaporated and the residue treated with water (50 ml) and then extracted with dichloromethane (3 x 20 ml). The combined dichloromethane extracts were dried (MgSO_4) and evaporated. The residue was fractionated by silica gel column chromatography with light petroleum (b.p. 40-60°C)-ethyl acetate, 2:1 as eluant giving (24S, 25S)-24-methyl-23-methoxyimino-25-phenyl-milbemycin X as a single isomer. ^{13}C n.m.r. (270 MHz; CDCl_3) included δ 155.43, 139.37, 128.32, 128.26, 127.61, 61.40, 40.63, and 11.01ppm.



The following compounds of formula (I), wherein $\text{R}_1 = \text{R}_3 = \text{H}$, were prepared using the methods of Examples 1 and 2. Throughout the Examples, the acetylenic precursors of formula $\text{HC} = \text{C}.\text{CR}^4\text{R}^5\text{CR}^6\text{R}^7\text{OP}$ were prepared using the routes described in EP-A-O 353 959 (USSN 387, 351):


Example 2A



Treatment of a methanolic solution of the title compound with 2M HCl for 16 h gave a small amount of the Z isomer which was separated from the E isomer by prep tlc (SiO_2 hexane/ethyl acetate (2:1)).



Table V




Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
1	1	OH	C ₆ H ₅	CH ₃	H	H	=O	TBDMS	A and B
2	2	OH	C ₆ H ₅	CH ₃	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N-} \\ \text{E isomer} \end{array}$	TBDMS	A and B
3	see below	OH	t-C ₄ H ₉	H	H	H	=O	TES TBDMS	A and B
4	1	OCH ₃			H	H	=O	TBDMS	B
5	1	OH			H	H	=O	TBDMS	B
6	see below	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N-} \\ \text{two isomers} \end{array}$	TBDMS	A and B
7	2	OH			H	H	$\begin{array}{c} \text{OMe} \\ \diagup \\ \text{=N-} \\ \text{E isomer} \end{array}$	TBDMS	B















Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
8	1	OH	Cyclo-C ₆ H ₁₁	CH ₃	H	H	=O	TBDMS	A
9	1	OH	H	H	tert-C ₄ H ₉	H	=O	TBDMS	A and B
10	2	OH	Cyclo-C ₆ H ₁₁	CH ₃	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
11	2	OH	t-C ₄ H ₉	CH ₃	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
12	1	OH	t-C ₄ H ₉	CH ₃	H	H	=O	TMS	A
13	1	OH		H	H	H	=O	TES	A
14	6	OH		H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	A


Ex. No	Prep. as per Ex. No	R2	R7	R4	R6	R5	R8	Acetylenic Precursor	
								Protecting group P	Route of preparation
15	6	OH		H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{Z isomer} \end{array}$	TES	A
16	2	OH	C ₆ H ₅	CH ₃	H	H	$\begin{array}{c} \text{NH} \\ \text{CONH}_2 \\ \text{E isomer} \end{array}$	TBDMS	A and B
17	2	OH	Cyclo-C ₆ H ₁₁	C ₂ H ₅	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
18	2	OH	C ₆ H ₅	$\text{p-C}_4\text{H}_9$	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
19	1	OH	4-CH ₃ .C ₆ H ₄	H	H	H	=O	TBDMS	A
20	6	OH	4-CH ₃ .C ₆ H ₄	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
21	6	OH	4-CH ₃ .C ₆ H ₄	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{Z isomer} \end{array}$	TBDMS	A

Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
22	2	OH	H	H	Cyclo-C ₆ H ₁₁	CH ₃	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	B
23	1	OH	H	H		H	=O	TBDMS	Commercial Compound
24	1	OH	Cyclo-C ₆ H ₁₁	H	H	H	=O	TBDMS	A
25	6	OH	Cyclo-C ₆ H ₁₁	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
26	6	OH	Cyclo-C ₆ H ₁₁	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{Z isomer} \end{array}$	TBDMS	A
27	1	OH		H	H	H	=O	TES	A
28	6	OH		H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	A

Ex. No	Prep. as per Ex. No	R2	R7	R4	R6	R5	R8	Acetylenic Precursor	
								protecting group P	Route of preparation
29	2	OH	H	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N} \\ \text{E:Z 1:1} \\ \text{mixture} \end{array}$	TBDMS	Commercial Compound
30	1	OH		H	H	H	=O	TES	A
31	6	OH		H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{=N} \\ \text{Z isomer} \end{array}$	TES	A
32	6	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OCH}_2\text{C}_6\text{H}_4\text{NO}_2 \\ \diagup \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	B
33	6	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OCH}_2\text{C}_6\text{H}_4\text{NO}_2 \\ \diagup \\ \text{=N} \\ \text{Z isomer} \end{array}$	TES	B
34	6	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OC}_2\text{H}_5 \\ \diagup \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	B
35	6	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OC}_2\text{H}_5 \\ \diagup \\ \text{=N} \\ \text{Z isomer} \end{array}$	TES	B

Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
36	see below	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OC}_4\text{H}_9-\text{t} \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	B
37	see below	OH	t-C ₄ H ₉	H	H	H	$\begin{array}{c} \text{OC}_4\text{H}_9-\text{t} \\ \text{=N} \\ \text{Z isomer} \end{array}$	TES	B
38	6	OH		H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E+Z isomers} \end{array}$	TES	A
39	6	OH	C ₄ H ₉ -t	CH ₃	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E+Z isomers} \end{array}$	TMS	A
40	6	OH	n-C ₆ H ₁₃	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TBDMS	A
41	6	OH	n-C ₆ H ₁₃	H	H	H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{Z isomer} \end{array}$	TBDMS	A
42	2	OH				H	$\begin{array}{c} \text{OCH}_3 \\ \text{=N} \\ \text{E isomer} \end{array}$	TES	B
43	6	OH	C ₄ H ₉ -t	H	H	H	$\begin{array}{c} \text{OH} \\ \text{=N} \\ \text{E+Z isomers} \end{array}$	TES	B

Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
44	6	OH	C ₄ H ₉ -t	H	H	H	 =N-  E isomer	TES	B
45	6	OH	C ₄ H ₉ -t	H	H	H	 =N-  Z isomer	TES	B
46	see below	OH	C ₄ H ₉ -t	H	H	H	-OH two epimers	-	-
47	6	OH	C ₄ H ₉ -t	H	H	H	 =N-  E+Z isomers	TES	B
48	6	OH	C ₄ H ₉ -t	H	H	H	 =N-  E+Z isomers	TES	B
49	6	OH	C ₄ H ₉ -t	H	H	H	 =N-  E+Z isomers	TES	B
50	2	OH	C ₄ H ₉ -t	H	H	H	 =N-  E+Z mixture	TES	B
51	2	OH	C ₄ H ₉ -t	H	H	H	 =N-  E+Z mixture	TES	B

Ex. No	Prep. as per Ex. No	R ²	R ⁷	R ⁴	R ⁶	R ⁵	R ⁸	Acetylenic Precursor	
								Protecting group P	Route of preparation
52	6	OH			H	H	=O	TES	B
53	54	OH		H	H	H	$\text{=N}-\text{OC}(\text{NHC}_6\text{H}_5)_2$ E + Z isomers mixture	TES	B
54	see below	OH		H	H	H	$\text{=N}-\text{OCH}_2\text{OCH}_3$ E + Z isomers	TES	B
55	6	OH		H	H	H	$\text{=N}-\text{O} \text{C}(\text{C}_2\text{H}_5)_3$ E+Z isomers	TES	B
56	see below	OH		H	H	H	$\text{=N}-\text{CH}_2\text{C}(\text{CH}_3)_3$ E+Z isomers	TES	B
57	see below	OH		H	H	H	$\text{=N}-\text{CH}_2\text{C}(\text{CH}_3)_3$ Two Epimers	TES	B

Example 3

5

23 Oxo-25(S)-t-butyl milbemycin x

To a solution of (4S)-5,5-dimethyl-4-triethoxysilyloxy-1-hexyne (8.3 g, 33 mmol) in THF (100 ml) at -78°C under a nitrogen atmosphere was added butyllithium (1.6M in hexane, 18.9 ml, 30 mmol) dropwise over a period of 5 mins. and the mixture stirred at -78°C for a further 3 h. A solution of VS 48927 (4.8g, 8.6 mmol) in THF (20 ml) was added to the mixture which was stirred at -78°C for a further 15 mins. The reaction was quenched with a cold solution (-20°C) of glacial acetic acid (10 ml) in THF (10 ml) and the mixture was then allowed to warm to 0°C. Brine (100 ml) was added and the mixture was extracted with ether (3 x 100 ml). The combined organic extracts were washed with water, dried (MgSO₄) and evaporated to an approximate volume of 50 ml. Methanol (50 ml) was added and the solution again evaporated to an approximate volume of 50 ml. 4-Toluenesulphonic acid (1g) was added and the mixture stirred at 20°C for 1 h.

Sodium bicarbonate (100 ml) was added to the mixture and the whole extracted with dichloromethane (3 x 100 ml). The combined organic extracts were washed with brine (100 ml), dried (MgSO₄) and evaporated. The residue was purified by column chromatography (silica eluted initially with dichloromethane and subsequently gradient eluted with 10-60% ethyl acetate in hexane) to afford the methylacetal (3.35g, 66%).

This product (3.35g 5.7 mmol) was dissolved in methanol (25 ml) and a solution of mercuric oxide (56 mg, 0.26 mmol) water (3.5 ml) and concentrated sulphuric acid (0.75 ml, 14 mmol) was added dropwise at 20°C. The mixture was stirred at 20°C for 2 h., water was added and the mixture was extracted with dichloromethane (3 x 100 ml). The combined organic layers were washed with sodium bicarbonate solution (100 ml), dried (MgSO₄) and evaporated to give the title ketone (3.2g) pure by nmr + tlc.

30

Example 3A23 Oxo-25(s)-t-butyl milbemycin X dimethyl acetal

35

Treatment of the ketone of example 3 with methanol and 4-toluenesulphonic acid for 24 h gave the title compound.

Example 6

23-Oxo-25(S)-t-butyl milbemycin X (50 mg, 0.09 mmol) was dissolved in methanol (5 ml). A solution of methoxylamine hydrochloride (50 mg, 0.60 mmol) in water (2 ml) was added and the mixture stirred at room temperature (1h). The reaction mixture was concentrated and then treated with water (30 ml) and extracted with ether (3 x 15 ml). the combined ethereal extracts were dried (MgSO₄) and evaporated. The 1:1 mixture of Z and E oximes was separated by silica gel preparative thin layer chromatography with hexane - ethyl acetate, 1:1 as eluant.

The 23(Z)-methoxyimino-25(S)-t-butyl milbemycin X was obtained as a white solid (yield 16 mg). M_z - (FAB Na⁺/Noba) 622 [MNa]⁺ 50% (relative intensity). Hplc retention time = 7.7 min.

The 23(E)-methoxyimino-25(S)-t-butyl milbemycin X was obtained as a white solid (yield 16 mg). M_z - (FAB Na⁺/Noba) 622 [MNa]⁺ 25% (relative intensity). Hplc retention time = 7.9 min.

Hplc conditions : Dynamax C18 column (25 cm x 4.6 mm id) eluted with methanol - water , 9 : 1 at 1 ml/min monitored at 245 nm.

55

Examples 36 and 37

* Ration is dependent upon the pH of the reaction mixture.

23 (E and Z)- t-butyloxyimino -25 (S)-t-butyl milbemycin X

To a solution of 23-keto-25(S)-t-butyl milbemycin X (60 mg 0.1 mmol) and sodium acetate (300 mg, 2.2 mmol) in methanol (3 ml) was added 0-t-butylhydroxylamine hydrochloride (50 mg, 0.4 mmol). The mixture was stirred at 20°C for 1 h., water (10 ml) was added and the whole mixture extracted with dichloromethane (3 x 15 ml). The combined organic extracts were washed with water, dried (MgSO₄) and evaporated to dryness. Purification by preparative t.l.c., (silica taper plate (Analtech^R) eluted with ethyl acetate/hexane 2:5) yielded the title oximes as a 4:1 mixture of E:Z oxime isomers (54 mg). Treatment of a portion of this mixture (30 mg) with methanol (2 ml) and hydrochloric acid (1M, 0.2 ml) gave a 1:1 E:Z ratio of oxime isomers. The two isomers were separated by preparative t.l.c. (silica taperplate eluted four times with chloroform).

23(Z)-t-butyloxyimino-25(S)-t-butyl milbemycin X: tlc R_f = 0.5 (Silica eluted three times with chloroform containing 1.5% ethanol) m/z (FAB Na⁺/Noba) (relative intensity) 664 [MNa]⁺ (95%).

23(E)-t-butyloxyimino-25(S)-t-butyl milbemycin X: tlc R_f = 0.45 (Silica eluted three times with chloroform containing 1.5% ethanol) m/z (FAB Na⁺/Noba) (relative intensity) 664 [MNa]⁺ (95%).

Hplc retention times : Dynamax 60A Silica column (25 cm x 4.6 mm id) eluted with chloroform/methanol 99:1 at 1 ml/min monitored at 245 nm.

Retention time 23-Z isomer 14.2 min.

Retention time 23-E isomer 15.4 min.

Example 4625 23(R)-hydroxy-25(S)-t-butyl-milbemycin X and 23(S) epimer

To a solution of 5-O-triisopropylsilyl-23-oxo-25(S)-t-butyl milbemycin X (50 mg, 0.069 mmol) (obtainable from the compound of Example 3) in methanol (5 ml) was added sodium borohydride (10 mg, 0.26 mmol) and the mixture was stirred at 20°C for 30 mins. Brine (25 ml) was added and the mixture was extracted with ether (3 x 10 ml). The combined ether extracts were washed with water, dried (MgSO₄) and evaporated to dryness. The residue was dissolved in methanol (3 ml), 4-toluenesulphonic acid (5 mg) added and the mixture was stirred at 20°C for 16 h. Saturated sodium bicarbonate solution (10 ml) was added and the mixture was extracted with dichloromethane (3 x 10 ml). The combined organic extracts were washed with water (10 ml), dried (MgSO₄) and evaporated to give the crude R and S alcohols. The two products were separated and purified using preparative t.l.c. taper plates. (Silica eluted with ethyl acetate/hexane 2:5)

1) 23(R)-hydroxy-25(S)-t-butyl-milbemycin X (20 mg) t.l.c. R_f = 0.45 (SiO₂/Ethyl acetate/hexane 1:1); m/z (FAB Na⁺/Noba) (relative intensity) 595 [MNa]⁺ (15%).

2) 23(S)-hydroxy-25(S)-t-butyl-milbemycin X (9 mg) t.l.c. R_f = 0.35 (SiO₂/Ethyl acetate/hexane 1:1); m/z (FAB Na⁺/Noba) (relative intensity) 595 [MNa]⁺ (10%).

Example 54(a) 5-Acetoxy-23-keto-25(S)-t-butyl milbemycin X

23-keto-25(S)-t-butyl milbemycin X was acetylated with acetic anhydride in pyridine in the normal manner. The resultant 5-acetoxy-23-keto-25(S)-t-butyl milbemycin X was purified by silica gel column chromatography with hexane-ethylacetate, 2:1 as eluant.

b) 5-Acetoxy-23-hydroxyimino-25(S)-t-butyl milbemycin X

A solution of hydroxylamine hydrochloride (0.34g, 4.9 mmol) in water (5 ml) was added to a mixture of 5-acetoxy-23-keto-25(S)-t-butyl milbemycin X (1.0g, 1.6 mmol), sodium acetate (0.78g, 5.7 mmol) and methanol (30 ml). The mixture was stirred at R.T. (45 min). Water (100 ml) was added and the mixture extracted with ether (3 x 70 ml). The combined ethereal extracts were dried (MgSO₄) and evaporated giving 5-acetoxy-23-hydroxyimino-25(S)-t-butyl milbemycin X (0.9 g; 88%).

(c) 23-(E and Z)-Methoxymethoxyimino-25(S)-t-butyl milbemycin X

3M -Methyl magnesium iodide (0.4 ml, 1.2 mmol) was added dropwise to a stirred solution of 5-acetoxy-23-hydroxyimino-25(S)-t-butyl milbemycin X (368 mg, 0.6 mmol) in HMPA (10 ml) under a nitrogen atmosphere. After 5 min. bromomethyl methyl ether (0.07 ml, 0.8 mmol) was added and the mixture stirred at R.T. (1h). Ether (40 ml) and 0.5 M -hydrochloric acid (50 ml) were added to the reaction mixture. The organic layer was washed with 0.5M -hydrochloric acid, water, saturated sodium chloride solution, dried (MgSO₄) and evaporated.

The residue, methanol (10ml) and 1M -sodium hydroxide (0.5ml) were stirred in ice (2h). Ethyl acetate (40ml) was added and the mixture washed with 0.5M -hydrochloric acid, water, dried (MgSO₄) and evaporated. Purification by silica gel column chromatography with hexane-ethyl acetate, 2:1 afforded a mixture of 23-(E and Z)-methoxymethoxyimino-25(S)-t-butyl milbemycin X. The E and Z isomers were separated by preparative t.l.c. with hexane - ethyl acetate, 1:1 as eluant.

Example 5623-(2,2-dimethylpropylimino)-25(S)-t-butyl milbemycin X

A solution of 23-keto-25(S)-t-butyl milbemycin X (60mg, 0.1mmol) and 2,2-dimethylpropylamine (0.2ml) in dry tetrahydrofuran (5ml) was stirred at 70°C in the presence of p-toluenesulphonic acid monohydrate (trace) and 4A molecular sieves for 16h. The cooled solution was filtered and concentrated to a brown solid. Separation by column chromatography on silica eluting with ethyl acetate:hexane (1:1 v/v) gave the title compound as a solid. Mass spectrum FAB (NoBA/Na) m/z 662 (MNa⁺).

Example 5723-(2,2-dimethylpropylamino)-25(S)-t-butyl milbemycin X

To a solution of 23-keto-25(S)-t-butyl milbemycin X (63mg, 0.11mmol) and 2,2-dimethylpropylamine (0.2ml) in methanol (2ml) at room temperature was added sodium cyanoborohydride (7mg, 0.11mmol) and 2N aqueous hydrochloric acid (0.1ml). The solution was stirred for 18h poured into water and extracted with ethyl acetate. The organic phase was dried (MgSO₄) and evaporated to a white solid. Separation by thin layer chromatography on silica eluting with ethyl acetate:hexane (7:3 v/v) gave both the 23(R)- and 23(S)-products. Mass spectrum FAB (NoBA-Na) m/z 664 (MNa⁺) for both products.

Reference Example (preparation of acetylenic precursor for Example 3)5,5-dimethyl-1-hexyn-4-ol

3,3-Dimethyl-1,2-epoxybutane (10g, 100 mmol) was slowly added (over 30 min) to a stirred mixture of lithium acetylide ethylene diamine complex (90%, 15.3g, 150 mmol) and dry DMSO (40 ml) under a nitrogen atmosphere. The temperature was maintained at 10-15°C throughout the addition and the reaction stirred at approx. 15°C for a further 3 h. The mixture was cooled in ice and cautiously treated with ice cold water (200 ml). The dark coloured solution was extracted with ether (3 x 50 ml) and the ether extracts dried (MgSO₄) and evaporated. The residue was purified by column chromatography (silica eluted with 40-60° PE/ether 9:1) to yield the alcohol (8.9g., 71%).

Resolution of 5,5-dimethyl-1-hexyn-4-ol(a) Preparation of (S)-O-acetylmandelate

A solution of (R,S)-5,5-dimethyl-1-hexyn-4-ol (15g, 119 mmol), (S)-O-acetylmandelic acid (23g, 119 mmol) and 4-dimethylaminopyridine (1.45g 11.9 mmol) in dichloromethane (300 ml) was cooled in ice and DCC (24.5g 119 mmole) added over a period of 45 mins. The resulting mixture was stirred at room temperature for 1 h. Tlc (Silica eluted with 20% ether in hexane) showed the reaction to be complete. The mixture was filtered and the filtrate washed successively with dilute hydrochloric acid, dilute sodium bicarbonate solution and brine. The extract was dried (MgSO₄) and evaporated. The resulting oil was dissolved in hexane (50 ml) and left to crystallise. Crystals of the (S)-O-acetylmandelate ester of (S)-5,5-dimethyl-1-hexyn-4-ol (13 g 72%) were recovered by filtration and washed with a small volume of hexane.

b) Hydrolysis of (S)-O-acetylmandelate


The (S)-O-acetylmandelate ester (13g) was dissolved in methanol (200 ml) and treated with a solution of potassium carbonate (30g) in water (50 ml). The mixture was stirred at room temperature for 5 days after which time tlc (silica eluted with 20% ether in hexane) indicated that the reaction was complete. Water (250 ml) was added and the mixture extracted with hexane. The hexane extract was washed with water and cautiously evaporated to remove most of the hexane. Purification of the residue by column chromatography (silica eluted with Petrol (40-60°)/ether (5:1) yielded the pure (S)-alcohol (5.2g 96%).

(S)-5,5-dimethyl-4-(triethylsilyloxy)-1-hexyne

A mixture of (S)-5,5-dimethyl-1-hexyn-4-ol (1.96g 16 mmol), triethylamine (4.5 ml, 32 mmol) and dry dichloromethane (30 ml) was cooled to 0°C under a nitrogen atmosphere. Triethylsilyl trifluoromethanesulphonate (5.4 ml, 24 mmol) was slowly added and the mixture stirred at 0°C for a further 40 min. The mixture was treated with saturated sodium bicarbonate solution (60 ml) and the organic layer separated. The aqueous layer was extracted with dichloromethane (3 x 20 ml) and the combined dichloromethane extracts dried (MgSO₄) and evaporated. The residue was purified by column chromatography (silica eluted with Petroleum ether 40-60°) to yield the title compound 3.55g (95%).


Claims

1. A compound of formula (I):




wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or optionally O-substituted oximino; R³ is hydrogen, optionally protected hydroxy, or a group 4-(α-L-oleandrosyl)-α-L-oleandrosyloxy or α-L-oleandrosyloxy wherein the terminal hydroxy group is optionally protected; R⁴, R⁵, R⁶ and R⁷ are the same or different and each is hydrogen or an organic radical; and R⁸ is an optionally substituted amino or imino group; with the proviso that the compound of formula (I) is not a compound of formula (E) or (F) above or a compound disclosed in EP-A-0 307 220.

2. A compound of formula (IA);




wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or optionally O-substituted oximino; R³ is hydrogen, optionally protected hydroxy, or a group 4-(α -L-oleandrosyl)- α -L-oleandrosyloxy or α -L-oleandrosyloxy wherein the terminal hydroxy group is optionally protected; R⁴, R⁵, R⁶ and R⁷ are the same or different and each is hydrogen or an organic radical; and R⁸ is optionally protected hydroxy or oxo, with the proviso that (a) when R³, R⁵ and R⁶ are hydrogen and R⁴ is methyl, then R⁷ does not have any of the values set out in Table III hereinabove, (b) when R³ is not hydrogen, R⁴ is methyl and R⁵ and R⁶ are hydrogen, then R⁷ does not have any of the values set out in Table IV hereinabove, (c) when R³, R⁵ and R⁶ are hydrogen and R⁴ is ethyl, then R⁷ is not (Z)-4-methylpent-2-en-2-yl, and (d) when R³ is not hydrogen, R⁴ is methyl, and R⁵ is hydrogen, the 25-position is not substituted in the same way as that of compound III of US-A-4,285,963, and (e) it is not a compound disclosed in EP-A-0 317 148, EP-A-0 307 220, EP-A-0 308 145, EP-A-0 350 187 or EP-A-0 335 541.

3. A compound selected from the group consisting of compounds of formula (IB):



wherein R², R⁴, R⁵, R⁶ and R⁷ and R⁸ have the values set out in Table V hereinabove.

4. A milbemycin or avermectin of partial formula (I):



wherein R⁴ to R⁸ have the values defined in claim 1 or claim 2.


5. A compound according to any one of claims 1 to 4, for use in the treatment or prophylaxis of endo- and ectoparasitic infestations of the human or animal body.

6. A method of eradicating arthropod or nematode infestations, which method comprises applying a compound according to any one of claims 1 to 4, or a derivative thereof to the arthropods or nematodes or to their environment.


7. A pesticidal composition comprising a compound according to any one of claims 1 to 4 or a derivative thereof together with an inert carrier or excipient.

8. A pharmaceutical or veterinary composition comprising a compound according to any one of claims 1 to 4, or a pharmaceutically or veterinarily acceptable derivative thereof together with a pharmaceutically or veterinarily acceptable carrier or excipient.


9. A process for the preparation of a compound of formula (I) as defined in claim 1 or a compound of formula (IA) as defined in claim 2, which process comprises (hydrating and) cyclizing a compound of formula (II), (IV) or (V):



(II)




(IV)




wherein R^1 to R^8 and R^{18} have the values set out in claims 1 and 2, and/or, optionally, converting a compound of formula (I) or (IA) to a different compound of formula (I) or (IA).

10. A process for the preparation of a compound of formula (I) as defined in claim 1, which process comprises (a) cyclizing a compound of formula (IV) as defined in claim 9 or (b) treating a corresponding compound of formula (1) wherein R^8 is oxo with an amino containing compound, and, optionally, converting the compound of formula (I) obtained to a different compound of formula (I).


11. A process for the preparation of a milbemycin or avermectin of partial formula (V):




wherein R^{14} to R^{17} are the same or different and each is selected from hydrogen and an organic radical; and R^{19} is optionally protected hydroxy, oxo, or an optionally substituted amino or imino group, which process comprises (hydrating and) cyclizing a corresponding milbemycin or avermectin of partial formula (vi), (vii) or (viii):



(vi)



(vii)




(viii)


wherein R^{14} to R^{19} have the values set out above; and, optionally, converting the compound of formula (v) obtained to a different compound of formula (v).

Claims for the following Contracting State: ES

1. A process for the preparation of a compound of formula (I):



wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or optionally O-substituted oximino; R³ is hydrogen, optionally protected hydroxy, or a group 4'-(α -L -oleandrosyl)- α -L -oleandrosyloxy or α -L -oleandrosyloxy wherein the terminal hydroxy group is optionally protected; R⁴, R⁵, R⁶ and R⁷ are the same or different and each is hydrogen or an organic radical; and R⁸ is an optionally substituted amino or imino group with the proviso that the compound of formula (I) is not a compound of formula (E) or (F) above or a compound disclosed in EP-A-0 307 220, or a compound of formula (IA):




wherein R¹ is hydrogen or optionally protected hydroxy; R² is alkoxy, optionally protected hydroxy, oxo or optionally O-substituted oximino; R³ is hydrogen, optionally protected hydroxy, or a group 4'-(α -L -oleandrosyl)- α -L -oleandrosyloxy or α -L -oleandrosyloxy wherein the terminal hydroxy group is optionally protected; R⁴, R⁵, R⁶ and R⁷ are the same or different and each is hydrogen or an organic radical; and R⁸ is optionally protected hydroxy or oxo, with the proviso that (a) when R³, R⁵ and R⁶ are hydrogen and R⁴ is methyl, then R⁷ does not have any of the values set out in Table III hereinabove, (b) when R³ is not hydrogen, R⁴ is methyl and R⁵ and R⁶ are hydrogen, then R⁷ does not have any of the values set out in Table IV hereinabove, (c) when R³, R⁵ and R⁶ are hydrogen and R⁴ is ethyl, then R⁷ is not (Z)-4-methylpent-2-en-2-yl, and (d) when R³ is not hydrogen, R⁴ is methyl, and R⁵ is hydrogen, the 25-position is not substituted in the same way as that of compound III of US-A-4,285,963, and (e) it is not a compound disclosed in EP-A-0 317 148 EP-A-0 307 220, EP-A 0 308 145, EP-A-0 350 187 or EP-A-0 335 541, which process comprises (hydrating and) cyclizing a compound of formula (II), (IV) or (V):

5

10

15




(II)

20

25


30



(IV)

35

40




(V)

45 wherein R^1 to R^6 and R^{18} have the values set out above and/or, optionally, converting a compound of formula (I) or (IA) to a different compound of formula (I) or (IA).

2. A process for the preparation of a compound of formula (I) as defined in claim 1, which process comprises (a) cyclizing a compound of formula (IV) as defined in claim 1 or (b) treating a corresponding compound of formula (I) wherein R^8 is oxo with an amino containing compound, and, optionally, converting the compound of formula (I) obtained to a different compound of formula (I).


3. A process for the preparation of a milbemycin or avermectin of partial formula (V):

55




(V)


wherein R^{14} to R^{17} are the same or different and each is selected from hydrogen and an organic radical; and R^{19} is optionally protected hydroxy, oxo, or an optionally substituted amino or imino group, which process comprises (hydrating and) cyclizing a corresponding milbemycin or avermectin of partial formula (vi), (vii) or (viii):



(vi)




(vii)



(viii)

wherein R^{14} to R^{19} have the values set out above; and, optionally, converting the compound of formula (v) obtained to a different compound of formula (v).


4. A process according to any preceding claim, for the preparation of a compound selected from the group consisting of compounds of formula (IB):



(IB)

wherein R^2 , R^4 , R^5 , R^6 and R^7 and R^8 have the values set out in Table V hereinabove.

5. A process according to any one of claims 1 to 3, for the preparation of a milbemycin or avermectin of partial formula (i):



(1)

wherein R^4 to R^8 have the values defined in claim 1.

6. Use of a compound of formula (I), (IA), (IB) or (i) in the manufacture of a medicament for use in the treatment of prophylaxis or endo- and ectoparasitic infestations of the human or animal body.

7. A method of eradicating arthropod or nematode infestations, which method comprises applying a compound of formula (I), (IA), (IB), or (i) or a derivative thereof to the arthropods or nematodes or to their environment.

8. A pesticidal composition comprising a compound of formula (I), (IA), (IB) or (i) or a derivative thereof together with an inert carrier or excipient.

9. A pharmaceutical or veterinary composition comprising a compound of formula (I), (IA), (IB) or (i) or a pharmaceutically or veterinarily acceptable derivative thereof together with a pharmaceutically or veterinarily acceptable carrier or excipient.



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 90305367.6
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.)
X	<u>EP - A2/A3 - 0 214 739</u> (MERCK & CO. INC.) * Claim 1; page 2 * ---	2,4-8	C 07 H 17/08 C 07 D 493/22 A 01 N 43/90 A 61 K 31/335
X	<u>EP - A1 - 0 274 871</u> (SANKYO COMPANY LIMITED) * Claim 1; page 5 * ---	2,4-8	
X	<u>EP - A2 - 0 246 739</u> (SANKYO COMPANY LIMITED) * Claims 1,17 * ---	2,4-8	
P,X	<u>EP - A2 - 0 340 932</u> (PFIZER LIMITED) * Claims 1,11 * ---	2,4-8	
D,A	<u>EP - A2 - 0 307 220</u> (AMERICAN CYANAMID COMPANY) * Claims 1,7,8 * ---	1-8	
D,A	<u>EP - A1 - 0 259 779</u> (AMERICAN CYANAMID COMPANY) * Claims 1,9,10 * ---	1-8	TECHNICAL FIELDS SEARCHED (Int. Cl.)
D,A	<u>EP - A2/A3 - 0 260 536</u> (AMERICAN CYANAMID COMPANY) * Claims 1,8,9 * ---	1-8	C 07 H 17/00 C 07 D 493/00
D,A	<u>EP - A1 - 0 260 537</u> (AMERICAN CYANAMID COMPANY) * Claims 1,10,11 * ---	1-8	
D,A	<u>EP - A2 - 0 308 145</u> (PFIZER LIMITED) * Claims 1,10 * -----	2,4-8	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 24-08-1990	Examiner PETROUSEK
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

